

Creation of the Universe

Victor N. Pervushin,
Bogoliubov Laboratory of Theoretical Physics,
Joint Institute for Nuclear Research, 141980 Dubna, Russia

2nd February 2008

Abstract

Possibilities for solution of the problem of creation of the Universe from a physical vacuum in the framework the General Relativity and modern quantum field theory are discussed in the context of the official doctrine accepted in Trinity College at the Newton time.

A Christians in Science Conference “Design and Purpose in the Universe”
Saturday 18th March 2006, St. Nicholas’ Church, Market Place, Durham, DH1 1JP "www.cis.org.uk"

Contents

1	Physical “Laws” and “Data”	2
2	Action and Interval as “Foundations of Physics”	2
3	A Frame for Constrained “Data”	2
4	Newton’s doctrine versus the “Trinity” one	3
4.1	Two “descriptions” of one relativistic object	3
4.2	Hilbert’s Geometro-Dynamics of the Universe	5
4.3	Relative Units	5
4.4	Electro-Weak Epoch versus the Planck one	6
5	Design and Purpose in the Universe	7
6	Discussion at the 42nd Karpacz Winter School of Theoretical Physics	11

1 Physical “Laws” and “Data”

The founder of the first physical theory Isaac Newton gave to it some pattern of the official doctrine accepted in his Trinity College. Beginning with Newton’s mechanics all physical theories remember to a some extent the Christian theory of a human soul being in two *kingdoms*: a *kingdom of laws* and a *kingdom of wills*.

In any physical theory, a *kingdom of laws* is a set of equations of motion obtained by varying an *action*. A *kingdom of wills* is associated with a set of initial data which Laplace still required for unambiguous solutions of the Newton equations in order “to explain the World without God”.

The initial data of an object are measured by a set of physical instruments and are given by a *will* of experimenters. *Laws* of nature do not depend on the initial data but results of application of these *laws* can depend. The Newton equation of a train does not depend on the initial data, i.e. on the *will* of a passenger of this train who chose his initial position and the speed of the train, but the final result of the solution of this equation is a consequence of both the *will* of the passenger and the *law* of nature.

In contrast to mathematician Laplace, physicist Newton understood that in any explanation of the Design and Purpose in the Universe physicists should take into account both the *kingdoms* of laws and “data” given by God in a comoving reference frame where the Universe was created.

This comoving frame is identified by modern observers with the Cosmic Microwave Background (CMB) radiation distinguished by its dipole component measured [1] in the frame of an Earth observer.

Formulations of Einstein’s General Relativity and quantum field theory in a concrete frame were fulfilled by the founders of modern physics (for details see [2] – [6])¹. Now I present here the results of our attempts with my collaborators [10] – [20] to use these formulations, in order to describe the creation of the Universe in the framework of modern physical theory in agreement with observational data.

2 Action and Interval as “Foundations of Physics”

The official doctrine accepted in Trinity College “*In the beginning was the Word ..., and the Word was God*” (John 1:1) can be consistent with the modern physics, if God can be visible due to His action. The “action” of the unified theory was restored by both experimentalists and theoreticians in the 19th and 20th centuries in the context of the field nature of matter and space-time in the form of a sum of Hilbert’s action of the Einstein General Relativity [23] and the action of the Weinberg - Salam - Glashow Standard Model of elementary particles [24] settled in the Riemannian space-time defined by a geometric interval. Scientists believe in that equations of motion obtained by varying this action and supplemented by the initial data in the CMB frame can explain the origin of the Universe.

3 A Frame for Constrained “Data”

In General Relativity (GR) and quantum field theories, a comoving frame was introduced by Dirac, Heisenberg and Pauli, Fermi, Schwinger, and other physicists until the 1960s on the level

¹I was an eye-witnesser of that at the beginning of the 1970s theoreticians restricted quantum field theory by describing mainly scattering processes that do not depend on the data together with their reference frames. This restriction allowed to formulate the frame free Faddeev – Popov method [21] that gave an essential simplification of the proof of renormalizability (finiteness) of the Standard Model marked by the Nobel Prize to t’Hooft and Veltman [22]. But this restriction lost the historical pathway of reference frames in physical theories beginning with Copernicus until Dirac [4] and Schwinger [5].

of the action (see [2] – [6]) because it was the most straight path to determine the spectrum of the elementary and collective excitations and to separate equations of motion from a set of *constraints* between the initial data. These *constraints* of the data in all modern relativistic theories are principally new elements in comparison with the Newton mechanics.

A contemporary colleague of Newton in Trinity College could find that the constrained relativistic systems led to the official Trinity doctrine rejected by Isaac Newton² in his dramatic discussion with the founders of this doctrine³.

4 Newton's doctrine versus the “Trinity” one

4.1 Two “descriptions” of one relativistic object

David Hilbert's formulation of GR⁴ in a comoving frame of reference can be an effective illustration of the adequateness of the Trinity doctrine to description of any relativistic systems (particle, string, universe in GR).

In order to demonstrate this adequateness, let us consider the problem of measurements of the life-time of an unstable particle being in a relativistic train moving with velocity 200 thousands km/s. If a Driver of the train measures its life-time 10 s, then a Pointsman at a railway station measures 14 s (see Fig. 1). What is a correct life-time, 10 s or 14 s?

Two different times of the same particle is contradiction, in the Newton doctrine⁵.

Two different times of the same particle are described by two different actions given in two different frames (comoving and rest), in the Einstein-type intermediate doctrine⁶, where the relation between these two times is only a kinematic one, but not a consequence of dynamic

²Introducing the standards of consistent mathematical proof into physics in terms of clear absolute notions Newton was looking for the sense of these notions in Arian theology, trying to get out of the Trinity College doctrine which preached relativism and logically inconsistent trinity. One can think that for Newton as a member of Trinity College at Cambridge University this situation was a paradox [25].

³One of the founders of the Trinity theology Basil the Great revealed limits of the *consistent Aristotle's logic* (see [26, 27]) which anticipated the famous Gödel theorems formulated in the 20th century for *formal arithmetical systems*: i. Any *consistent* description of a system is not *complete*,

ii. Any *complete* description of a system is not *consistent* [28].

Here *consistent* means the absence of logical contradictions. The theological deed of Basil the Great was the positive constructive generalization of the *consistent Aristotle's logic*, in contrast with the negative Gödel theorems. Basil's generalization of *Aristotle's logic* can be formulated as a continuation of the first two Gödel theorems:

iii. Any *complete* description of a system with logical contradictions becomes *consistent* in a new meaning, if there are simultaneously two (or more) descriptions of the system connected by *relationships*, so that two opposite assertions about the same system belong to two their different *descriptions*.

⁴In the year of celebration of the 90th anniversary of GR we can now distinguish Einstein's treatment of general coordinate transformations (who considered them as generalization of the frame transformations [29]) from Hilbert's variation approach to GR [23], where the general coordinate transformations are considered as *diffeomorphisms* of the GR action and a *geometric interval*. There is an essential difference between the frame group of the Lorentz – Poincaré-type [30] leading to a set of initial data and the *diffeomorphism group* of general coordinate transformations restricting these initial data by *constraints*. This difference was revealed by two Nöther theorems [31]. The formulation of GR in terms of the Fock simplex [32] defined as a diffeo-invariant Lorentz vector helps us to separate diffeomorphisms from transformations of frames of reference.

⁵Resolving the Newton equation (the acceleration is equal to zero) one can find a trajectory of the train, i.e. the time dependence of the coordinate of the train $X(t) = X_I + V_I t$, where V_I is a ratio of the momentum P and a mass m . The passenger can choose another frame with other data, but the time of a particle in all frames will be unique and absolute, as it was postulated by Newton in his mechanics with the energy $E_{\text{Newton}}(P) = P^2/(2m)$.

⁶Lorentz, Poincaré, and Einstein saw that electrodynamics was consistent with mechanics, where the train had the trajectory $X(X_0) = X_I + \frac{P}{E(P)} X_0$, and the energy $E(P) = \sqrt{m^2 + P^2}$. This energy revealed and explained nuclear energetics. If an observer chooses another frame, the time X_0 converts into another one mixing with the coordinate by the kinematic Lorentz transformation in accordance with the Maxwell electrodynamics. This fact of changing time is one of relativistic effects meaning that each frame has its own time of a particle.




Figure 1: At the top of Fig.1 a relativistic train is depicted with an unstable particle. The life-time of this particle is measured by two observers, by a Pointsman and a Driver. Each of the observers has his time: Pointsman - variable X^0 , and Driver - geometrical interval η (and his wave function). At the bottom of the figure there is an image of the universe where each observer has two sets of measurable quantities corresponding to two observers of the particle. To the Pointsman there corresponds a field set of measurable quantities (mass φ scaled by the cosmological scale factor and density of a number of particles), and the Driver – geometrical set of measurable quantities (time interval η and initial data) [14, 15, 17].

equations of motion.

Two different times of the same particle and their relation as a consequence of dynamic equations of motion are described by one Hilbert's geometro-dynamic action given in one concrete frame, where there is a bifurcation of the classical time into two different times (time-variable measured by a Pointsman and time-interval measured by a Driver) of the same object. Isaac Newton could know in this bifurcation the Trinity doctrine, because these two times belong to two different *descriptions* (here the dynamic and the geometric) of the object by the same action in the same frame⁷.

The Hilbert geometro-dynamics of a particle leads not only to Einstein-type theory with the energy explaining nuclear processes, but also predicts an antiparticle with a negative energy known as a Dirac positron and reveals the historical pathway of quantum field theory (the main tool of high energy physics in the 20th century) because the quantum field theory is nothing, but primary and secondary quantizations of the energy constraint, in order to remove a negative energy and to make a particle stable. However, the quantum field theory uses only a time-variable measured by an external observer (i.e. a Pointsman) being out of a particle.

⁷In order to give the dynamic description of the relativistic relations, one should consider the time X_0 as one of the dynamic variables in the World space of events $[X_0|X_1]$ on equal footing with spatial coordinate $X = X_1$ and introduce one more time as a geometric measure η of a trajectory $X_0(\eta) = \frac{P_0}{m}\eta$, $X_1(\eta) = X_{1I} + \frac{P_1}{m}\eta$ of a particle in the World space of events $[X_0|X_1]$. These equations are supplemented by the energy constraint $P_0^2 - E^2 = 0$, $E = \pm\sqrt{P_1^2 + m^2}$ as an equation of the metric component of the Einstein interval expressed in terms of an unmeasurable “coordinate time” as an object of “general coordinate transformation” in the one-dimensional Riemannian space.

4.2 Hilbert's Geometro-Dynamics of the Universe

In the Universe, an observer is simultaneously a Driver and a Pointsman. To the Pointsman there corresponds a field set of measurable quantities (the cosmological scale factor⁸ considered as a *time-like variable* and densities of matter), and the Driver – geometrical set of measurable quantities (*time-interval* and initial data). In this case, geometro-dynamic relation between the *time-variable* and *time-interval*⁹ is just the famous Hubble law: the further a star, the more redshift of the star photons. As the cosmological scale factor is the *time-variable*, its momentum is the *energy of the Universe*¹⁰ in the field space of events. In such the Einstein – Hilbert GR the cosmic evolution is consistent with primary and secondary quantizations of the energy constraint¹¹ justified by the experience of the high energy physics the 20th century. These quantizations gave the description of the creation of the Universe and its geometric time and matter from the physical vacuum¹² [14, 15, 18, 20] in satisfactory agreement with observational data, if we chose the relative units [13].

4.3 Relative Units

The cosmic dynamics of the Universe was revealed in General Relativity by Alexander Friedmann [35] who kept in the Einstein interval only a cosmological scale factor and resolved equations of motion in this case. However, the interpretation of this cosmic dynamics as the expansion of the universe is possible only in the Newton doctrine of absolute units, if we propose that our standards belong to the kingdom of *laws*.

In accordance with the Trinity doctrine [26, 27], a man is free to choose his standard himself, because we can cognize only a ratio of things. If a measurement is one of the tools of cognition, the Trinity doctrine means that we can measure any physical quantity only in units of another physical quantity accepted as a standard¹³.

Defining a measurable interval of the length as the ratio of a Friedmann-like interval to the standard one that also belongs to Friedmann-like intervals, one can see that the measurable interval of the length is not expanding, as it does not depend on the cosmological scale factor.

Really, the scale factor disappears from the relative measurable interval, but not from the equations of motion. The equations of motion contain the Friedmann cosmological scale factor as a measure of all masses. It was shown [13, 14, 15, 17] that the relative units give a completely different physical picture of the evolution of the universe than the absolute units of the standard

⁸In GR, the cosmic evolution is extracted by a scale transformation $g_{\mu\nu} = a^2 \tilde{g}_{\mu\nu}$ well known as the cosmological perturbation theory [33, 34]. The GR action in terms of metrics $\tilde{g}$ depends on only the running Planck $\varphi(\eta) = a M_{\text{Planck}} \sqrt{3/8\pi}$ mass scaling all masses.

⁹The metric component $\sqrt{-\tilde{g}} \tilde{g}^{00} = (\tilde{N}_d)^{-1}$ gives the conformal time of photons flying in the conformal flat space-time $\zeta_{(\pm)} = \int dx^0 \langle N_d^{-1} \rangle^{-1} = \pm \int_{\varphi^0}^{\varphi} d\varphi / \langle (\tilde{T}_0^0)^{1/2} \rangle$, where $\langle F \rangle = V_0^{-1} \int d^3x F$ is the averaging over finite volume $V_0 = \int d^3x$, and $\tilde{T}_0^0$ is the energy-momentum tensor component.

¹⁰The energy constraint takes the form $P_\varphi^2 - E_\varphi^2 = 0$, where P_φ is the scale factor canonical momentum and $E_\varphi = 2 \int d^3x (\tilde{T}_0^0)^{1/2}$ should be treated as the “frame energy” of the Universe like $E = mc^2$ in Special Relativity.

¹¹The primary quantization of the energy constraint $[\hat{P}_\varphi^2 - E_\varphi^2] \Psi_L = 0$ leads to the unique wave function Ψ_L of the collective cosmic motion.

¹²The secondary quantization $\Psi_L = [1/\sqrt{2E_\varphi}][A^+ + A^-]$ and Bogoliubov's transformation: $A^+ = \alpha B^+ + \beta^* B^-$ diagonalizes the equations of motion by the condensation of “universes” $\langle 0|(i/2)[A^+ A^+ - A^- A^-]|0 \rangle = R$ and describes creation of a “number” of universes $\langle 0|A^+ A^-|0 \rangle = N$ from the stable Bogoliubov vacuum $B^-|0 \rangle = 0$.

¹³Maxwell revealed that the description of results of experimental measurement of electromagnetic phenomena by the field theory equations depends on the definition of measurable quantities in the theory and the choice of their measurement standard. In the introduction to his “A Treatise on Electricity and Magnetism” Maxwell wrote: “The most important aspect of any phenomenon from mathematical point of view is that of a measurable quantity. I shall therefore consider electrical phenomena chiefly with a view to their measurement, describing the methods of measurement, and defining the standards on which they depend.”[36].




Figure 2: The Hubble diagram [13] in cases of the absolute units of standard cosmology (SC) and the relative ones of conformal cosmology (CC). The points include 42 high-redshift Type Ia supernovae [37] and the reported farthest supernova SN1997ff [38]. The best fit to these data requires the dominance of the cosmological constant density (70%) and 25% of the Cold Dark Matter density, in the case of the Friedmann standard cosmology, whereas for the relative units these data are consistent with the dominance of density of a free scalar field ($85\% \pm 10\%$) with the square root dependence of the scale factor on conformal time (that corresponds to the stiff state) [13, 39].

cosmology. The spectrum of photons emitted by atoms from distant stars billion years ago remains unchanged during the propagation and is determined by the mass of the constituents at the moment of emission. When this spectrum is compared with the spectrum of similar atoms on the Earth which, at the present time, have larger masses, then a redshift is obtained.

The temperature history of the expanding universe copied in the relative units looks like the history of evolution of masses of elementary particles in the cold universe with a constant temperature of the cosmic microwave background. The relative observable distance loses the cosmological scale factor a , in comparison with the absolute one. Therefore, in this case, the observational redshift – coordinate-distance relation [37, 38] corresponds to the dominance of the stiff state of free scalar field with the square root dependence of the scale factor on conformal time [13] (see also Fig. 2). Just this time dependence of the scale factor on the measurable time (here – conformal one) is used for description of the primordial nucleosynthesis [40].

4.4 Electro-Weak Epoch versus the Planck one

Thus, the relative units doctrine leads to a single stiff state for all epochs including the creation of a quantum universe at the beginning. In terms of the relative units the Planck mass loses its status as the fundamental parameter of the equations of motion and becomes the present-day value of the running mass scale in contrast with the Inflationary Model [34, 41] based on the proposal about the existence of the Planck epoch at the beginning. The initial data of the creation are determined by parameters of matter cosmologically created from the stable quantum vacuum at the beginning of a universe. In the Standard Model, W-, Z-vector bosons have maximal




Figure 3: The surface shows the distribution function of creation of longitudinal ($N_Z(x)$) components of the W-bosons in units of the dimensionless time ($\tau = 2\eta H_I$) and the dimensionless momentum ($x = q/M_I$) with the constant conformal temperature $T_c \sim (M_I^2 H_I)^{1/3} = (M_W^2 H_0)^{1/3} \sim 3K$. These bosons created from vacuum at the moment when their Compton length defined by the inverse mass $M_I^{-1} = (a_I M_W)^{-1}$ is close to the universe horizon defined in the stiff state as $H_I^{-1} = a_I^2 (H_0)^{-1}$. Equating these quantities $M_I = H_I$ one can estimate the initial data of the scale factor $a_I^2 = (H_0/M_W)^{2/3} = 10^{-29}$ and the Hubble parameter $H_I = 10^{29} H_0 \sim 1 \text{ mm}^{-1} \sim 3K$. CMB radiation is described as the product of decay of primordial W-,Z- bosons during the time-life $\eta_L \sim (2/\alpha_W)^{2/3} (T_c)^{-1}$ expressed in terms of the Weinberg coupling constant $\alpha_W = \alpha_{\text{QED}}/\sin^2 \theta_{\text{Weinberg}} \sim 0.03$ [15, 17]. The primordial mesons before their decays polarize the Dirac fermion vacuum and give the baryon asymmetry frozen by the CP – violation so that for billion photons there is only one baryon, and relative contributions of the baryon matter and radiation are in agreement with observational data: $\Omega_b \sim \alpha_W \sim 0.03$, and $\Omega_R \sim 10^{-5} \div 10^{-4}$ [17]. The equations of the longitudinal vector bosons in SM, in this case, describe the “power primordial spectrum” of the CMB radiation.

probability of this cosmological creation due to their mass singularity at the moment when their Compton length is close to the universe horizon defined by the primordial Hubble parameter. Therefore, the Universe was a factory of W-,Z-vector bosons at the beginning. The observational data on CMB reflect the parameters of Standard Model of elementary particles (see Fig. 3,4) [15, 17]¹⁴.

The geometro-dynamic formulation of General Relativity [2, 4, 23, 32] supplemented by the cosmological scale factor as a collective time-like variable in a comoving frame and “relative” units [17, 18, 20] corresponds rather the Trinity doctrine than the Newton one. This geometrodynamics allows us not only to restore the historical pathway of physical description of the Universe evolution but also to use experience of quantum field theory formulation verified by the high energy physics experiments. We gave here a set of numerous arguments in favor that this quantum unified theory (GR & SM & an additional scalar field) can be a theoretical basis of the further detailed investigation of astrophysical observational data including CMB fluctuations as one of the highlights of present-day cosmology with far-reaching implications and more precise observations are planned for the near future.

5 Design and Purpose in the Universe

Thus, the papers of the founders of modern physics [2, 4, 23, 29, 32] allow us to restore the historical pathway of description of the Universe evolution in the comoving frame with the

¹⁴Now it is a single possibility to explain the CMB observations [1] in the framework of GR and SM, as it was shown in [18, 20] the standard cosmological perturbation theory [33, 34] applied for analysis of the CMB observations [1] in the Inflationary Model [41] has a different number of variables than the Einstein General Relativity [29].




Figure 4: The curve shows power law of the temperature angular power spectrum with first peak $l \sim 210$ [1]. The parameters of CMB can reflect parameters of the SM of elementary particles, in particular the Weinberg coupling constant [15, 17, 20].

definite initial data in the unified theory like the description of the trajectory of a relativistic particle in SR after its quantum creation from stable vacuum. This description reveals the trinity of times in GR: i) the unmeasurable coordinate time in Hilbert's action as an object of general coordinate transformations (i.e., diffeomorphisms), ii) the diffeo-invariant geometric time-interval measured by our watch, and iii) the time-like variable as the cosmological scale factor measured by astrophysical observations and considered as a measure of the Universe evolution in the field space of events, where the Universe was created [11] – [20].

One can say that modern physics goes beyond the bounds of the Newton Arian doctrine accepted by modern theoreticians [42] who have a possibility to continue the Newton dialogue with the founders of the Trinity doctrine [26, 27]. Who is a man in the Universe? He is a passenger who lost his rest frame in the “frame free” method, who uses only absolute units and parameters of the “natural” laws including the “absolute Planck mass”, and cannot determine unambiguously the energy of the Universe? Or he is a Driver, who knows his frame and place, chooses his relative units, measures the initial data, introduces new concepts and changes his logics as one of the tools of his cognition, who is able to determine the Universe energetics, describes its creation, and is responsible for its further fate, as he has his free “will”, in order to take part in further creation of the measurable world and to have an eternal life of his unmeasurable soul in the kingdom of His will?

References

- [1] Spergel D.N., *et al.* “First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Determination of Cosmological Parameters”, *Asrophys. J. Suppl.* **148**, 175–203 (2003); astro-ph/0302209.
- [2] Dirac P.A.M. “Generalized Hamiltonian Dynamics”, *Proc. Roy. Soc. (London)* **A 246**, 326–332 (1958); Dirac P.A.M. “Fixation of Coordinates in the Hamiltonian Theory of Gravitation”, *Phys. Rev.* **114**, 924–930 (1959).

- [3] Arnowitt R., Deser S., and Misner C.W. “Canonical Variables for General Relativity”, *Phys. Rev.* **117**, 1595–1602 (1960).
- [4] Dirac P.A.M. “Quantum Theory of Emission and Absorption of Radiation”, *Proc. Roy. Soc.*, **A 114**, 243–249 (1927);
Dirac P.A.M. “Gauge Invariant Formulation of Quantum Electrodynamics”, *Can. J. Phys.*, **33**, 650–661 (1955).
- [5] Schwinger J. “Non-Abelian Gauge Fields. Relativistic Invariance”, *Phys. Rev.*, **127**, 324–330 (1962).
- [6] Polubarinov I. V. “Equations of Quantum Electrodynamics”, *Physics of Particles and Nuclei*, **34**, 377–410 (2003).
- [7] Pervushin V.N. “On the Vacuum in the Gauge Theories”, *Riv. del Nuovo Cimento* **8**, N 10, 1–48 (1985).
- [8] Nguyen Suan Han and Pervushin V. N. “Operator Foundation of Functional Integral in Non-Abelian Gauge Theory”, *Mod.Phys.Lett.* **A2**, 367–371 (1987).
- [9] Pervushin V.N. “Hadron QCD (Bound States in Gauge Theories)”, *Nucl.Phys. B (Proc. Supp.)* **15**, 197–206 (1990); Pervushin V. N. and Nguyen Suan Han “Hadronization and Confinement in QCD”, *Can.J.Phys.* **69**, 684–691 (1991); Preprint CERN TH 5866/90, Geneve, 1990; Blaschke D., Kalinovsky Yu.L., Munchow L., Pervushin V.N., Ropke G., and Schmidt S. “Instantaneous Chiral Quark Model for Relativistic Mesons in a Hot and Dense Medium”, *Nucl.Phys.* **A 586**, 711–733 (1995).
- [10] Gogilidze S., Ilieva Nevena, Pervushin V.N. “Dirac variables and zero modes of Gauss constraint in finite volume two -dimensional QED”, *Int. J. Mod. Phys.* **A 14**, 3531–3542 (1999); Pervushin V. N. “Dirac Variables in Gauge Theories”, *Physics of Particles and Nuclei* **34**, 348–376 (2003); *Lecture Notes in DAAD Summerschool on Dense Matter in Particle - and Astrophysics*, JINR, Dubna, Russia, August 20 - 31, 2001; [hep- th/ 0109218]; Lantsman L.D. and Pervushin V.N. “Monopole Vacuum in Non-Abelian Theories”, *Physics of Atomic Nuclei* **66**, 1384–1394 (2003).
- [11] Gyngazov L.N., Pawlowski M., Pervushin V.N., and Smirichinski V.I. “Proper Time Dynamics in General Relativity and Conformal Unified Theory”, *General Relativity and Gravitation* **37**, 128–38 (1998).
- [12] Pawlowski M., Papoyan V. V., Pervushin V. N., Smirichinski V.I. “Conformal Unification of General Relativity and Standard Model for Strong and Electroweak Interactions”, *Phys.Lett.* **B 418**, 263–269 (1998).
- [13] Behnke Danilo, Blaschke David B., Pervushin Victor N., and Proskurin Denis “Description of Supernova Data in Conformal Cosmology without Cosmological Constant”, *Phys. Lett.* **B 530**, 20–26 (2002); [gr-qc/0102039].
- [14] Barbashov B.M., Pervushin V.N., and Proskurin D.V. “An Excursus into Modern Cosmology”, *Physics of Particles and Nuclei* **34**, Suppl. 1, S68–S90 (2003).
- [15] Pervushin V.N. “Early Universe as a W-, Z- Factory”, *Acta Physica Slovakia* **53**, 237–243 (2003).
- [16] Gusev A., Flin P., Pervushin V., Vinitsky S., and Zorin A., “The Universe evolution as a possible mechanism of formation of galaxies and their clusters”, *Astrophysics* **47** 242–246 (2004); [astro-ph/0301543]; Flin P., Pervushin V., and Zorin A., “Capture of cosmic objects by central gravitational field of a galaxy cluster”, astro-ph/0406051.
- [17] Blaschke D.B., Vinitsky S.I., Gusev A.A., Pervushin V.N., and Proskurin D.V. “Cosmological Production of Vector Bosons and Cosmic Microwave Background Radiation”, *Physics of Atomic Nuclei* **67**, 1050–1062 (2004); [hep-ph/0504225].
- [18] Barbashov B.M., Zakharov A.F., Zinchuk V.A., and Pervushin V.N. “Hamiltonian General Relativity in Finite Space and Cosmological Potential Perturbations”, submitted in *Int.J.Mod.Phys.* **A**; astro-ph/0511824.
- [19] Zakharov A.F., Zinchuk V.A., and Pervushin V.N. “Tetrad Formalism and Frames of References in General Relativity”, *Physics of Particles and Nuclei* **37** 104–134 (2006).

- [20] Barbashov B.M., Pervushin V.N., Zakharov A.F., and Zinchuk V.A., “Hamiltonian Cosmological Perturbation Theory”, *Phys. Lett. B* **633** 458–462 (2006); [hep-th/0501242].
- [21] Faddeev L. and Popov V. “Feynman diagrams for the Yang-Mills field”, *Phys. Lett. B* **25**, 29–35, (1967).
- [22] ‘t Hooft G. and Veltman M. “Regularization and renormalization of gauge fields”, *Nucl. Phys.*, **B 44**, 189–2001 (1972).
- [23] Hilbert D. “Die Gründlangen der Physik”, *Nachrichten von der Kön. Ges. der Wissenschaften zu Göttingen, Math.-Phys. Kl.*, **3**, 395–407 (1915).
- [24] Bardin D. and Passarino G. “The standard model in the making: precision study of the electroweak interactions”, Clarendon, Oxford, 1999.
- [25] Manuel F.E. “The religion of Isaac Newton”, Oxford, 1974;
Westfall R.S. “Never at rest: A biography of Isaac Newton”, Cambridge, 1982;
Koyré A. “Newtonian studies”, L., 1965.
- [26] Florovsky G.V. “Orient fathers of the 4th century”, Paris, 1931; Russian translation, Moscow, 1992.
- [27] Meyendorff John “Byzantyne Theology. Trends and Doctrinal Themes”, Fordham University Press, N.Y., 1979.
- [28] Gödel K. “Über formal unentscheidbare Satze der Principia Mathematica und verwandter Systeme I”, *Monatshefte für Math. und Phys.* **38**, 173–198 (1931); (Reprinted and translated in Godel, Kurt, 1986, Collected Works, **1**, 144–195, Oxford: Oxford University Press).
- [29] Einstein A. “Die Gründlange der allgemeinen Relativitätstheorie”, *Ann.d.Phys.* **49**, 769–826 (1916).
- [30] Poincare H. “Sur la dynamique de l’electron”, *Rendiconti del Circolo matematico di Palermo* **21** 129 (1906).
- [31] Nöther E., *Invariante Variationsprobleme Nachrichten von der Kön. Ges. der Wissenschaften zu Göttinger, Math.-Phys. Kl.*, **2** 235–258 (1918).
- [32] Fock V.A. “Geometrisierung der Diracschen Theorie des Electrons”, *Zs. Phys.* **57**, 261–277 (1929).
- [33] Lifshits E. M. “On the gravitational stability of the expanding universe”, *ZhETF* **16**, 587–602 (1946).
- [34] Mukhanov V. F., Feldman H. A., and Brandenberger R. H. “Theory of cosmological perturbations”, *Phys. Rep.* **215**, 203–333 (1992).
- [35] Friedmann A. A., “Über die Krümmung des Raumes”, *Z. für Phys.* **10** 377–386 (1922) (English translation in: *Gen. Rel. Grav.* **31**, 1991–2000 (1999)); Friedmann A. A., “Über die Möglichkeit einer Welt mit konstanter negativer Krümmung des Raumes”, *Z. für Phys.* **21** 326–332 (1924) (English translation in: *Gen. Rel. Grav.* **31**, 2001–2008 (1999)).
- [36] Maxwell J.C. “A Treatise on Electricity and Magnetism”, Clarendon Press, Oxford, 1873.
- [37] Perlmutter, S., et al. “Measurements of Omega and Lambda from 42 High-Redshift Supernovae”, *Astrophys. J.* **517**, 565–586 (1999); [astro-ph/9812133].
- [38] Riess A. G., et al. “Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant”, *Astron. J.* **116**, 1009–1038 (1998); [astro-ph/9805201].
- [39] Behnke D. “Conformal Cosmology Approach to the Problem of Dark Energy”, PhD Thesis, Rostock Report MPG-VT-UR 248/04 (2004).
- [40] Weinberg S. “First Three Minutes. A modern View of the Origin of the universe”, Basic Books, Inc., Publishers, New-York, 1977.
- [41] Linde A.D. “Elementary Particle Physics and Inflation Cosmology”, Nauka, Moscow, 1990.
- [42] Duff M.J., “The world in eleven dimensions: supergravity, supermembranes and M-theory”, I.O.P. Publishing 1999, <http://bookmark.iop.org/bookpage.htm/bookB5>.

6 Discussion at the 42nd Karpacz Winter School of Theoretical Physics

I am very grateful to Profs. A. Borowiec and M. Francaviglia for the hospitality at the 42nd Karpacz Winter School of Theoretical Physics on Current Mathematical Topics in Gravitation and Cosmology, Łądek, Poland, 6-11.02.2006 (<http://www.ift.uni.wroc.pl/karp42/>). At the School there were very interesting and sharp debates on energy in General relativity (GR), its variables, and a reference frame of the Universe creation. I list here some questions and my responses.

FRAME?

Question by Prof. Zbigniew Oziewicz (Universidad Nacional Autonoma de México):

What does “the concrete reference frame of the Universe creation” mean? A frame is the coordinate basis, and physical results should not depend on any “basis”.

Response (V.P.):

Let me recall that the introduction of the physical concept of a “reference frame” as a three-dimensional coordinate basis with a watch was really stimulated by the great Polish astronomer M. Kopernik who proposed in 1543 the heliocentric “frame” as an alternative to the Ptolemaeus “frame” connected with the Earth observer.

Thus, there appeared a definition of “physics” as a science about observations (later – measurements) of objects being in the “comoving frame” of their center of masses by devices associated with the “rest frame” of an observer. In the context this definition Kopernik’s “relativity” is nothing, but different epicycles “relative” to different frames.

Galilei in his “Dialog ...” in 1632 converted these two frames into a set of inertial frames moving with constant velocities. He identified one of the inertial frames with a ship, where a motion of the measurable object “relative” to a moving device is described by the difference of their two velocities. In this case, you are right, different motions of the ship do not change indications of the devices, i.e. initial data. I meant only motions of a comoving frame relative to a rest frame, i.e. transformations of the initial data, because we now have a similar situation with the Cosmic Microwave Background (CMB) measured by the SkyLab “Hubble” devices being in the “Ptolemaeus (rest) frame” moving “relative” to CMB with the velocity 400 km/s to Leo. If we move the SkyLab “Hubble” together with CMB (i.e. make the Lorentz transformations changing the initial data), the dipole components of CMB temperature disappears (see [19]). Thus, in accordance with Kopernik’s “relativity” principle, the Universe as any “observable” object has its “comoving frame” marked by CMB.

Question by Prof. Zbigniew Oziewicz:

What is your opinion about the relation of the Lorentz group to electrodynamics?

Response (V.P.):

The Faraday – Maxwell electrodynamics contained the hidden symmetries revealed later by Poincare (known as the Lorentz group of transformations of the initial data) and by Weyl (known as gauge symmetry leading to constraints of initial data). Einstein’s “Special Relativity” is his generalization of mechanics based on the Lorentz group of frame transformations leading to different times “relative” to different frames and energy $E = mc^2$ in the comoving frame.

Einstein’s “General Relativity” was treated by him as generalization of the Lorentz frame transformations; whereas Hilbert in his “Foundations” [23] considered GR as a “gauge” theory with a set of constraints revealed by the Hilbert theorem known as the second Nöther one. The Lorentz transformations were introduced in GR by means of the Fock tetrads [32].

VARIABLES?

Problem of the choice of variables in different theories of gravitation was considered in Lectures by Profs. **M. Francaviglia**, **G. Allemandi** (both from University of Torino) and **L. Sokołowski** (Jagiellonian University, Kraków). In all these lectures the remarkable fact was noted. All metric theories of gravitations are connected by general conformal transformations of their variables and they are mathematically equivalent. The questions appeared: What variables can be considered as measurable quantities? and What is the criterion of the choice of a true gravitation theory among the mathematically equivalent ones?

Response (V.P.)

In the context of the results presented in these Lectures the action of Einstein's GR is mathematically equivalent to the negative action of the conformal invariant theory of a scalar field (called "dilaton") [12, 13]. The latter does not contain any dimensional fundamental parameter of the type of the Planck mass. If this action is supplemented by the SM one, where the Higgs mass is replaced by the "dilaton" field, we get the unified theory, where the scale (and, therefore, conformal) symmetry can be broken by the initial data at the beginning of the Universe provided that the present day value of the "dilaton" field is equal to the Planck mass. We see that the idea of conformal symmetry of the World developed by Richard Rączka (see cites in [12]) excludes the Planck epoch at the beginning of the Universe. Conformal symmetry distinguishes the Lichnerowicz "conformal variables" $g_{\mu\nu}^{(L)} = |g^{(3)}|^{-1/3} g_{\mu\nu}$ constructed with the help of the spatial metric determinant and used by Dirac in his Hamiltonian approach [2]. Identification of the Lichnerowicz "conformal variables" with the measurable quantities leads to Conformal Cosmology with varying masses (instead of the expanding Universe) where Supernova Data are described by the ordinary homogeneous free scalar field in the stiff state without Cosmological Constant [13].

ENERGY?

The heaviest debates at the Winter School were about the choice of the energy in General Relativity and cosmology. Profs. **J. Garecki** (University of Szczecin), **L. Lusanna** (INFN, Firenze), **L. Sokołowski**, and **M. Dyrda** (both from Jagiellonian University, Kraków) in their talks used the conventional definition of the ADM-type energy relative to an external observer in asymptotically flat infinite spacetime. Profs. **M. Francaviglia** (University of Torino), **J. Lukierski** (ITF, University of Wrocław) and other participants insisted on that just this definition could not be applied in cosmology because an internal observer has a possibility to observe only finite spacetime of the Universe. The illustration of this discussion was the remarkable film by **Lorenzi Marcella** (Cosenza, Italy) about the Einstein energy $E = mc^2$.

Response (V.P.)

In the Hilbert-type formulation of Special Relativity (SR), $E = mc^2$ is considered as a solution of the energy constraint with respect to the canonical momentum of the time-like variable. Just the GR generalization of this SR energy $E = mc^2$ to the Universe in finite space was the topic of my talk [11, 18, 19, 20], where the collective time-like variable is identified with the homogeneous scale factor a multiplied by the Planck mass. It is a varying Planck mass $\varphi = aM_{\text{Pl}}\sqrt{3/(8\pi)}$ considered as the homogeneous "dilaton" degree of freedom in conformal version of GR. The Einstein-type energy in GR (as a solution of the energy constraint) depends on the varying mass φ . Just this dependence leads to the freedom to create the Universe and its matter from the vacuum in the comoving frame with the initial data $\varphi_I^2/\varphi_0^2 = a_I^2 = H_0/H_I \sim 10^{-29}$ [15, 17].

Question by Prof. M. Szydłowski (Jagiellonian University, Kraków)

What does creation of universes from the vacuum mean?

Response (V.P.):

QFT of universes is formulated by analogy with the QFT of particles [4] as the primary and the secondary quantizations of the energy constraint, in the context of Hilbert's variational principle [23], in order to remove the negative Einstein-type energy and to make stable the quantum system¹⁵. We need only to understand: What does QFT mean for the “varying masses”? But this problem was solved at the end of the 60s for particles by Chernikov and Parker with the help of the Bogoliubov transformation (1946) and for universes where the role of the “mass” is played by the cosmological energy density in 2006 [18, 19, 20].

DEGREES OF FREEDOM?

Question by Prof. S. Odintsov (Barcelona University & Tomsk University)

What is the status of the conformal sector of Quantum Gravity considered in the papers

I. Antoniadis, E. Mottola, PRD V. 45, 1992, p.2013;

S.D. Odintsov, Zeit.fur Physik V. C54, 1992, p.531;

I. Antoniadis and S.D. Odintsov, Phys.Lett. B343, 1995, p.76 ?

Response (V.P.):

These papers were devoted to GR in infinite spacetime without any supposition about a stable vacuum, negative energy, initial data, time-variable, units of measurement, and separation of all variables and metric components into degrees of freedom with gauge-invariant initial data violating the frame symmetry and potentials without initial data, and they explained nothing.

In particular, the kinetic energy density of the conformal sector is negative. Therefore, Dirac (in his Hamiltonian approach to GR formulated in the “comoving frame” [2]) identified all scalar components of the conformal sector with potentials. This identification is in agreement with the Schwarzschild solution, but not with cosmological evolution that appears in the conformal sector, in the case of finite spacetime, as the homogeneous zero mode of the momentum constraint, i.e the scale factor. Lifshits [33] treated this cosmological scale factor as an additional homogeneous variable. However, this treatment leads to double counting of the homogeneous variable and loses a possibility to formulate a Hamiltonian approach together with quantization and the freedom to create the Universe from the vacuum [19].

Dirac had no homogeneous degree of freedom (with Hamiltonian and without cosmology), and Lifshits introduced two (with cosmology and without Hamiltonian).

In the finite volume Hamiltonian approach to GR [19, 20] the scale factor is introduced as the spatial averaging the spatial determinant logarithm¹⁶. The negative sign of the energy density of the cosmological scale factor shows us that it is the evolution parameter, and the constraint value of its canonical momentum is the constrained energy like the Einstein one $E = mc^2$ in SR.

¹⁵The existence of a stable vacuum as the state with minimal energy can be considered as the quantum analog of the Boltzmann H-theorem about the existence of stable temperature.

¹⁶In this case the Newton potential is defined in the class of function with nonzero Fourier harmonics, i.e., the spatial averaging of this potential is equal to zero. This fact reveals a difference of the ADM-type energy based on this potential from the cosmological energy density given in finite space by means of the spatial averaging.

COSMOLOGY?

The set of Lectures by **Salvatore Capozziello** (University of Napoli) was devoted to the analysis of “an impressive amount of different astrophysical data” in the class of the Λ - Dark Matter cosmological models with the present-day accelerating expansion, where the nature of the Λ dark energy is still unknown. In this class of models, all principal cosmological problems (homogeneity, flatness, horizon, etc) are solved by the inflation mechanism [41]. The main test of Inflationary Model is the derivation of “CMB primordial power spectrum” [34] based on the cosmological perturbation theory [33]. There were a lot questions at the School: What is difference of the finite volume Hamiltonian GR [20] from the Inflationary Model [34, 41]?

Response (V.P.)

The main differences between the finite volume Dirac Hamiltonian approach to GR [20] and the Inflationary Model (IM) are the following.

1. In the finite volume GR the observational quantities of “the cosmological scale factor” and the “conformal time” are identified with the spatial volume averaging the metric scalar components, so that their cosmological perturbations do not contain the zero Fourier harmonics, in contrast to the Lifshits cosmological perturbation theory [33] used in IM [34, 41], where the “conformal time” is determined by the physically unattainable procedure of synchronization of watches in the whole Universe, and “the cosmological scale factor” is introduced as an additional variable that increases the number of variables.

2. In the finite volume Hamiltonian GR the Dirac principle of stability of a relativistic quantum system is used in contrast with IM. The Dirac “stability” means i) the zero canonical momentum fields are gauge-invariant potentials, ii) existence of the vacuum as a state with a minimal constrained energy, iii) the zero momentum of the local volume element removing its negative energy density and leading to a nonzero shift-vector N^i of the coordinate origin with spacial oscillations¹⁷ and the potential scalar perturbations, whereas the Lifshits-type kinetic scalar perturbations explaining CMB spectrum in IM disappear.

3. The CMB spectrum can be reproduced, if the absolute units of length in IM are replaced by the relative units for which SN data are compatible with the state of Dark Energy as a free scalar field [15]. In this case, the Early Universe is a factory of W-,Z- bosons [17] the equations of which reproduce the Lifshits-type kinetic scalar perturbations explaining CMB spectrum in IM. This factory gave us all matter and CMB as the decay product of the primordial bosons created from the stable “vacuum” after the creation of the Universe together with its coordinate frame and time interval known here as the Kopernik comoving frame.

In contrast to the IM in which the cosmic evolution begins with the Planck epoch and the Planck mass is treated as a fundamental parameter of the “law”, in the Hamiltonian GR the Planck epoch becomes the present-day one and the Planck mass is treated as the law free datum in the comoving frame of the Universe.

RESUME

It is useful to remember that Newton in his Trinity College strongly distinguished the *kingdom of laws* from the *kingdom of freedom (of will)*. Newton knew that any *law* does not depend on *will*, but application of the *law* does. The creation of the Universe is the application of the *law*. The instruction of this application can present here as the historical pathway of physics: from empiric facts to laws and from *laws to the grace* (John 1:17) of freedom to create the Universe from “nothing” (Newton called this history the *shoulders of giants*):

¹⁷The test of the finite volume Dirac Hamiltonian GR is the large-scale structure of the Universe (see Fig. 5).

Ptolemaeus' rest frame (RF),
 Kopernik's comoving frame (CF),
 Galilei's inertial frames as a set of all possible initial data given without the law (i.e. free),
 Newton's differential law invariant with respect to transformations of the law free initial data,
 Faraday's program: the field nature of matter, and unification of forces of Nature,
 Maxwell's equations with hidden Lorentz's frame symmetry (Poincare) and gauge one (Weyl),
 Einstein's Special Relativity (SR) as different times for different frames (RF & CF),
 Einstein's General Relativity (GR) as generalization of the Lorentz frame group?,
 Hilbert's variational principle for GR, where Einstein's generalization becomes gauge group,
 Fock's tetrad as the separation of the Lorentz frame transformations from the gauge ones,
 Friedmann revelation of the cosmological scale factor as the collective motion in GR,
 Einstein's $E_{\pm} = \pm mc^2$ in Hilbert's constrained version of SR supplemented by the interval,
 Dirac QFT for particles as the primary and secondary quantization of the energy constraint,
 Lichnerowicz variables $g_{\mu\nu}^{(L)} = |g^{(3)}|^{-1/3} g_{\mu\nu}$ distinguished by the spatial metric determinant,
 Dirac - ADM Hamiltonian approach to GR in terms of radiation-type Lichnerowicz variables,
 Zel'manov's global gauge symmetry of the Hamiltonian GR as foundation of cosmic evolution,
 Finite volume Dirac Hamiltonian approach to GR with the cosmic motion and its energy

$E_{\varphi} = 2 \int d^3x^{(L)} \sqrt{T_0^{0(L)}}$ as the canonical momentum of $\varphi = aM_{\text{Pl}} \sqrt{3/(8\pi)}$ [20],

Hubble law in the conformal units as the spatial averaging the Lichnerowicz interval,

Primary and secondary quantization of the energy constraint by Bogoliubov's transformation,

Dirac – Schwinger “fundamental operator quantization” [5] of Weinberg – Salam – Glashow Standard Model in the Kopernik-type comoving frame of the Universe, where all field components are separated into “potentials” (without initial data) and “degrees of freedom” (of initial data) with different poles of their propagators.

Recall that Faddeev proved the theorem of equivalence of “fundamental operator quantization” [5, 7] with the frame free “Lorentz gauge formulation” (where all field components are considered as “degrees of freedom” on equal footing) only for the scattering amplitudes. Nobody proved even in QED that instantaneous atoms and molecules formed by the Coulomb potential can be obtained by the Faddeev-Popov heuristic quantization [21] in the frame free Lorentz gauge where all propagators have only the light cone singularities. To his great surprise a contemporary theoretician can know that Schwinger “*rejected all Lorentz gauge formulations as unsuited to the role of providing the fundamental operator quantization*” [5]. Thereupon, it is worth emphasizing that Schwinger following Dirac postulated the higher priority of the quantum principles in comparison with relativistic ones [2, 4]¹⁸.

¹⁸Modern M-theories with D=26, D=10,11 [42] are consequences of the “scattering branch” of the Faddeev – Popov-type heuristic quantization [21]. The fundamental operator quantization of a relativistic string leads to R  hrlich spectrum without the Virasoro algebra [18]. In any case, the SN data [38, 37] in relative units compatible with Conformal Dark Energy is described by a scalar field, so that the scalar sector of the sum of GR and SM can be the conformal D=6 brane in the 4th-dimensional spacetime.




Figure 5: The diffusion of a system of particles moving in the space $ds^2 = d\eta^2 - (dx^i + N^i d\eta)^2$ with periodic shift vector N^i and zero momenta could be understood from an analysis of the following system of O.D.E. $dx^i/d\eta = N^i$ considered for the two-dimensional case. For the system we reproduce the results of numerical simulations given by Klyatskin V.I. “Stochastic Equations” M. Fizmatlit, 2001, where $t = \eta m_{(-)}, \overline{m}_{(-)} = H_0 \sqrt{6[\Omega_R(z+1)^2 + (9/2)\Omega_{\text{Matter}}(z+1)]/7}$ [20]. The size ~ 130 Mpc of spatial oscillations of matter is in agreement with value of radiation-type density $\Omega_R \sim 3 \cdot 10^{-3}$ at the time of diffusion $z_r \sim 1100$ [20].